



## Emeraid® Piscivore

### Product Description

Emeraid Piscivore is the first elemental diet designed for the critically ill seabird.

### Recommended Use

This diet may be fed as a sole or partial source of nutrition.

After months of feeding Emeraid Piscivore to hundreds of seabirds from dozens of species, International Bird Rescue has recommended Emeraid Piscivore as a superior diet in the stabilization of debilitated seabirds.

### Features and Benefits

Emeraid Piscivore is a fish-oil based diet with fat levels that can be modified from 5% to 30%. Fish oil is added to Emeraid powder in-house.

This diet contains:

- All essential amino acids, making protein more easily absorbable
- Hydrolyzed protein and purified amino acids
- Stabilized vitamins
- Dietary nucleotides to provide DNA and RNA precursors
- Chelated minerals

### Product Sizes

400 g tub

2 kg tub

### Guaranteed Analyses

Crude protein (min) 44.50% Crude fat (min) 4.40% Crude fiber (max) 4.46% Moisture (max) 9.00% Metabolizable energy 4.55 kcal/g Omega-3 fatty acids (min) 0.34% Omega-6 fatty acids (min) 2.86% Calcium (min) 0.96% Phosphorus (min) 0.84%

### Caloric Distribution

Protein 22.3 % Fat 30.1 % Digestible carbohydrates 47.8 %

## Storage and Shelf Life

- Refrigerate Emeraid after opening.
- Refrigerate or freeze fish oil.

## Preparation and Administration

Tube fed formula is absorbed best when the bird is warm and hydrated. In the adult bird, crop or stomach volume may be estimated as 50 ml/kg. In the debilitated patient or one with a past history of regurgitation, administer one-third to one-half of estimated stomach volume for the first feeding. Increase the volume fed over two to three feedings.

1. Formula should ideally be prepared immediately before feeding and no more than 30 minutes before feeding. Reheating is not advised.
2. Measure three level large scoop of powder (equivalent to 18.86 grams). Do not pack.
3. Mix in very warm water 105-110°F (41-43°C) with fish oil just before feeding.
4. Add 90-180 ml of water per 1 scoop of powder.
5. Dilute mixture 50% with water for dehydrated birds.
6. Stir mixture vigorously using a whisk or mixing ball.
7. Reduce the volume of Emeraid formula fed as the bird's appetite for fish improves.

## Ingredients

Soy protein hydrolyzed, corn syrup solids, cellulose, defatted wheat germ, corn oil, glutamine (an essential amino acid), dicalcium phosphate, ground limestone, potassium chloride, DL-methionine (an essential amino acid), magnesium sulfate, vitamin A supplement, vitamin E supplement, vitamin D3 supplement, menadione sodium bisulfite complex (source of vitamin K activity), choline chloride, cysteine (an essential amino acid), taurine (an essential amino acid), ethoxyquin (a preservative), manganese oxide, niacin supplement, calcium dipantothenate, riboflavin supplement, thiamine mononitrate, pyridoxine hydrochloride, vitamin B12 supplement, folic acid, zinc oxide, copper sulfate, biotin, ascorbic acid, sodium selenite.

## Distribution for European Zoos:

[www.kiezebrink.eu](http://www.kiezebrink.eu)

